A budget of paradoxes

—

The beginnings and aims of Hilbert's school

András Máté

9th October 2026

Russell paradox in naïve set theory

Russell paradox in naïve set theory

We have unlimited comprehension:

$$\exists y \forall x (x \in y \leftrightarrow A(x))$$

for any open sentence $A(x)$.

Russell paradox in naïve set theory

We have unlimited comprehension:

$$\exists y \forall x (x \in y \leftrightarrow A(x))$$

for any open sentence $A(x)$.
Therefore,

$$\exists y \forall x (x \in y \leftrightarrow x \notin x)$$

Russell paradox in naïve set theory

We have unlimited comprehension:

$$\exists y \forall x (x \in y \leftrightarrow A(x))$$

for any open sentence $A(x)$.
Therefore,

$$\exists y \forall x (x \in y \leftrightarrow x \notin x)$$

Let r be a such y (existential instantiation), and let us substitute r for x, too (universal instantiation).

$$r \in r \leftrightarrow r \notin r$$

Russell paradox in naïve set theory

We have unlimited comprehension:

$$\exists y \forall x (x \in y \leftrightarrow A(x))$$

for any open sentence $A(x)$.
Therefore,

$$\exists y \forall x (x \in y \leftrightarrow x \notin x)$$

Let r be a such y (existential instantiation), and let us substitute r for x, too (universal instantiation).

$$r \in r \leftrightarrow r \notin r$$

We have proved a logical falsity from the (unlimited) comprehension using only logical rules.

An embarrassing analogy

An embarrassing analogy

Cantor's theorem: For any set, there is no one-to-tone correspondence between the set and its power set.

An embarrassing analogy

Cantor's theorem: For any set, there is no one-to-tone correspondence between the set and its power set.
Be H any set, $\mathcal{P}(H)$ its power set, and f an injective mapping from H to $\mathcal{P}(H)$. We show that there is at least one member of $\mathcal{P}(H)$ that is not in the range of f:

An embarrassing analogy

Cantor's theorem: For any set, there is no one-to-tone correspondence between the set and its power set.
Be H any set, $\mathcal{P}(H)$ its power set, and f an injective mapping from H to $\mathcal{P}(H)$. We show that there is at least one member of $\mathcal{P}(H)$ that is not in the range of f:

$$H_0 =: \{x : x \notin f(x)\}$$

An embarrassing analogy

Cantor's theorem: For any set, there is no one-to-tone correspondence between the set and its power set.
Be H any set, $\mathcal{P}(H)$ its power set, and f an injective mapping from H to $\mathcal{P}(H)$. We show that there is at least one member of $\mathcal{P}(H)$ that is not in the range of f:

$$H_0 =: \{x : x \notin f(x)\}$$

Suppose (for contradiction) that $H_0 = f(h)$.

$$h \in f(h) \leftrightarrow h \notin f(h)$$

Russell's paradox in Frege's *Grundgesetze* system

Russell's paradox in Frege's *Grundgesetze* system

Let us consider the concept R: 'to be the value range of a concept that is false for its own value range'.

Russell's paradox in Frege's *Grundgesetze* system

Let us consider the concept R: 'to be the value range of a concept that is false for its own value range'.

Formally, $R(x) \leftrightarrow_{def} \exists G(x = {}^{\vee}G \wedge \neg G(x))$.

${}^{\vee}F$ is the value range of the function F.

Russell's paradox in Frege's *Grundgesetze* system

Let us consider the concept R: 'to be the value range of a concept that is false for its own value range'.

Formally, $R(x) \leftrightarrow_{def} \exists G(x = {}^{\backprime\backprime}G \wedge \neg G(x))$.

${}^{\backprime\backprime}F$ is the value range of the function F.

Let us substitute ${}^{\backprime\backprime}R$ for x.

$$R({}^{\backprime\backprime}R) \leftrightarrow \exists F({}^{\backprime\backprime}R = {}^{\backprime\backprime}F \wedge \neg F({}^{\backprime\backprime}R))$$

Russell's paradox in Frege's *Grundgesetze* system

Let us consider the concept R: 'to be the value range of a concept that is false for its own value range'.

Formally, $R(x) \leftrightarrow_{def} \exists G(x = {}^{\backprime\backprime}G \wedge \neg G(x))$.

${}^{\backprime\backprime}F$ is the value range of the function F.

Let us substitute ${}^{\backprime\backprime}R$ for x.

$$R({}^{\backprime\backprime}R) \leftrightarrow \exists F({}^{\backprime\backprime}R = {}^{\backprime\backprime}F \wedge \neg F({}^{\backprime\backprime}R))$$

Because of the first conjunct in the scope of $\exists$, any concept F which makes the existential quantification true is true for the same objects as R (by Axiom V). Therefore, the right side is true iff $\neg R({}^{\backprime\backprime}R)$.

The central problem: paradoxes

The central problem: paradoxes

Russell's paradox: published in Russell's *Principles of Mathematics* (1903)

The central problem: paradoxes

Russell's paradox: published in Russell's *Principles of Mathematics* (1903)

Frege immediately notes that Cantor's set theory contains precisely the the same inconsistency.

The central problem: paradoxes

Russell's paradox: published in Russell's *Principles of Mathematics* (1903)

Frege immediately notes that Cantor's set theory contains precisely the the same inconsistency.

Another paradox in set theory (Burali-Forti) becomes known some years earlier.

The central problem: paradoxes

Russell's paradox: published in Russell's *Principles of Mathematics* (1903)

Frege immediately notes that Cantor's set theory contains precisely the the same inconsistency.

Another paradox in set theory (Burali-Forti) becomes known some years earlier.

The central theme of foundational research/philosophy of mathematics: how can paradoxes be eliminated paradoxes and how can we avoid tha recurrence of such problems?

The central problem: paradoxes

Russell's paradox: published in Russell's *Principles of Mathematics* (1903)

Frege immediately notes that Cantor's set theory contains precisely the the same inconsistency.

Another paradox in set theory (Burali-Forti) becomes known some years earlier.

The central theme of foundational research/philosophy of mathematics: how can paradoxes be eliminated paradoxes and how can we avoid tha recurrence of such problems?

A collection of relevant paradoxes follows (with reference to De Morgan (1872) *A budget of paradoxes*).

The Liar paradox

The Liar paradox

(L) The sentence in the first line of this frame is false.

The Liar paradox

(L) The sentence in the first line of this frame is false.

If the sentence L is true, then its content holds, therefore L – that is the sentence in the first line – is false.

The Liar paradox

(L) The sentence in the first line of this frame is false.

If the sentence L is true, then its content holds, therefore L – that is the sentence in the first line – is false.

If L is false, then the sentence that claims that L is false is true, therefore L is true.

The Liar paradox

(L) The sentence in the first line of this frame is false.

If the sentence L is true, then its content holds, therefore L – that is the sentence in the first line – is false.

If L is false, then the sentence that claims that L is false is true, therefore L is true.

$$L \leftrightarrow \neg L$$

Variants for the Liar

Variants for the Liar

Liar-circle:

$p_1 \leftrightarrow \neg p_2$, $p_2 \leftrightarrow \neg p_3$, $\ldots$ $p_{2n-1} \leftrightarrow \neg p_1$

Variants for the Liar

Liar-circle:

$p_1 \leftrightarrow \neg p_2,\ p_2 \leftrightarrow \neg p_3,\ \ldots\ p_{2n-1} \leftrightarrow \neg p_1$

$p_1 \leftrightarrow \neg p_2,\ p_2 \leftrightarrow \neg p_3\ ,\ \ldots p_{2n} \leftrightarrow p_1$

Variants for the Liar

Liar-circle:

$p_1 \leftrightarrow \neg p_2$, $p_2 \leftrightarrow \neg p_3$, $\ldots$ $p_{2n-1} \leftrightarrow \neg p_1$

$p_1 \leftrightarrow \neg p_2$, $p_2 \leftrightarrow \neg p_3$, $\ldots$ $p_{2n} \leftrightarrow p_1$

Strenghtened Liar:
Let us allow that 'is false' and 'is not true' are not the same. That is, there are sentences that are neither true nor false („gappy").

Variants for the Liar

Liar-circle:

$p_1 \leftrightarrow \neg p_2$, $p_2 \leftrightarrow \neg p_3$, $\ldots$ $p_{2n-1} \leftrightarrow \neg p_1$

$p_1 \leftrightarrow \neg p_2$, $p_2 \leftrightarrow \neg p_3$, $\ldots$ $p_{2n} \leftrightarrow p_1$

Strenghtened Liar:
Let us allow that 'is false' and 'is not true' are not the same. That is, there are sentences that are neither true nor false („gappy").

$$L_S \leftrightarrow (L_S \text{ is not true})$$

Burali-Forti paradox

Burali-Forti paradox

Let BF the class of all ordinals, well-ordered by the relation $<$ (i.e., $\in$).

Burali-Forti paradox

Let BF the class of all ordinals, well-ordered by the relation $<$ (i.e., $\in$).

It is an ordinal. It is larger than any ordinal because any ordinal is a member of it.

Burali-Forti paradox

Let BF the class of all ordinals, well-ordered by the relation $<$ (i.e., $\in$).

It is an ordinal. It is larger than any ordinal because any ordinal is a member of it.

It is smaller than its successor.

Two more famous paradoxes

Two more famous paradoxes

Let us call a one-place predicate F heterological iff $F(F)$ is false. E. g. 'abstract' is abstract, but 'red' is not red. Is 'heterological' heterological?

Two more famous paradoxes

Let us call a one-place predicate F heterological iff $F(F)$ is false. E. g. 'abstract' is abstract, but 'red' is not red. Is 'heterological' heterological?
Known as Grelling-Nelson, Weyl, or simply heterological-paradox.

Two more famous paradoxes

The smallest number that cannot be defined in English in less than 81 characters

Richard's paradox

Richard's paradox

There are countably many real numbers between 0 and 1 that can be defined by a finitely long definition.

Richard's paradox

There are countably many real numbers between 0 and 1 that can be defined by a finitely long definition.

Let us enumerate all these numbers in the sequence a_k. Consider the following real number:

$$a = 0.d_1 d_2 ... d_n ...,$$

where $d_n = 6$ if the nth digit after the decimal point of of a_n is 5 and $d_n = 5$ otherwise.

Richard's paradox

There are countably many real numbers between 0 and 1 that can be defined by a finitely long definition.

Let us enumerate all these numbers in the sequence a_k. Consider the following real number:

$$a = 0.d_1 d_2 ... d_n ...,$$

where $d_n = 6$ if the nth digit after the decimal point of of a_n is 5 and $d_n = 5$ otherwise.

a differs from any member of our sequence, but it is defined.

The test and the hypergame

The test and the hypergame

The teacher says: Next week you'll have a test. I won't tell you which day. You will be surprised.

The test and the hypergame

The teacher says: Next week you'll have a test. I won't tell you which day. You will be surprised.

Students take the test on Wednesday and are surprised.

The test and the hypergame

G is an ordinary game between two players iff it ends in finitely many moves. H is the following hypergame: the first player chooses an ordinary game, and then they play it. Is H an ordinary game or not?

Russell's vicious circle principle

Russell's vicious circle principle

'Whatever involves *all* of a collection must not be one of the collection;' or, conversely: 'If, provided a certain collection had a total, it would have members only definable in terms of that total, then the said collection has no total.'

'Mathematical logic as based on the theory of types', 1908

Russell's vicious circle principle

'Whatever involves *all* of a collection must not be one of the collection;' or, conversely: 'If, provided a certain collection had a total, it would have members only definable in terms of that total, then the said collection has no total.'

'Mathematical logic as based on the theory of types', 1908

Self-reference: a sentence refers to itself, i.e. its truth conditions contain some condition on its truth or falsity.

Russell's vicious circle principle

'Whatever involves *all* of a collection must not be one of the collection;' or, conversely: 'If, provided a certain collection had a total, it would have members only definable in terms of that total, then the said collection has no total.'

'Mathematical logic as based on the theory of types', 1908

Self-reference: a sentence refers to itself, i.e. its truth conditions contain some condition on its truth or falsity.

Or it contains a quantification over propositions including the proposition expressed by the sentence.

Russell's vicious circle principle

'Whatever involves *all* of a collection must not be one of the collection;' or, conversely: 'If, provided a certain collection had a total, it would have members only definable in terms of that total, then the said collection has no total.'

'Mathematical logic as based on the theory of types', 1908

Self-reference: a sentence refers to itself, i.e. its truth conditions contain some condition on its truth or falsity.

Or it contains a quantification over propositions including the proposition expressed by the sentence.

Russell's principle (called vicious circle principle) prohibits self-reference. It seems sufficient to avoid the previous paradoxes.

Yablo's paradox

Yablo's paradox

Let us consider the following infinite sequence $p_1, p_2, ..., p_n, ...$ of propositions:

$$p_n \leftrightarrow \forall k(k > n \rightarrow \neg p_k)$$

Yablo's paradox

Let us consider the following infinite sequence $p_1, p_2, ..., p_n, ...$ of propositions:

$$p_n \leftrightarrow \forall k(k > n \rightarrow \neg p_k)$$

Stephen Yablo, 1989

Yablo's paradox

Let us consider the following infinite sequence $p_1, p_2, ..., p_n, ...$ of propositions:

$$p_n \leftrightarrow \forall k(k > n \rightarrow \neg p_k)$$

Stephen Yablo, 1989

It is a liar-like, but infinitary paradox that does not violate the vicious circle principle and does not contain any sort of self-reference.

Aims of foundational research

Aims of foundational research

- Create firmly–based/infallible theories

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency
- Create rich (possibly omniscient) theories

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency
- Create rich (possibly omniscient) theories

Three ways out of the trap of paradoxes:

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency
- Create rich (possibly omniscient) theories

Three ways out of the trap of paradoxes:

1. Improve logic and produce a unique general theory free of risks (logicism)

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency
- Create rich (possibly omniscient) theories

Three ways out of the trap of paradoxes:

1. Improve logic and produce a unique general theory free of risks (logicism)
2. Risky theories but a reliable metatheory (formalism)

Aims of foundational research

- Create firmly–based/infallible theories
- Avoid inconsistency
- Create rich (possibly omniscient) theories

Three ways out of the trap of paradoxes:

1. Improve logic and produce a unique general theory free of risks (logicism)
2. Risky theories but a reliable metatheory (formalism)
3. Abandon the priority of logic in favor of a more reliable basis (intuitionism)

Hilbert's foundational program at the beginnings

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid
- Change of perspective: not the science of the space, but the science of anything that satisfies the axioms

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid
- Change of perspective: not the science of the space, but the science of anything that satisfies the axioms

Another programmatic claim: Existence in mathematics is nothing but consistency.

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid
- Change of perspective: not the science of the space, but the science of anything that satisfies the axioms

Another programmatic claim: Existence in mathematics is nothing but consistency.

Ideal case: categorical theories

A theory is categorical iff all its models are isomorphic.

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid
- Change of perspective: not the science of the space, but the science of anything that satisfies the axioms

Another programmatic claim: Existence in mathematics is nothing but consistency.

Ideal case: categorical theories

A theory is categorical iff all its models are isomorphic.

Metamathematics is the (new) branch of mathematics investigating theories as mathematical objects (with the aim to prove that they are consistent, complete, categorical, etc.).

Hilbert's foundational program at the beginnings

Hilbert (*Foundations of Geometry*, 1st edition 1899): Axioms implicitly define the basic notions of the theory.

- Improved version of Euclid
- Change of perspective: not the science of the space, but the science of anything that satisfies the axioms

Another programmatic claim: Existence in mathematics is nothing but consistency.

Ideal case: categorical theories

A theory is categorical iff all its models are isomorphic.

Metamathematics is the (new) branch of mathematics investigating theories as mathematical objects (with the aim to prove that they are consistent, complete, categorical, etc.).

Standard way to investigate theories in the metatheory: formalize them in first-order logic.

Formalism, Hilbert's program, Hilbert's school

Formalism, Hilbert's program, Hilbert's school

Formalism: a way of philosophizing about mathematics – older than the *Grundlagenkrise*.

Formalism, Hilbert's program, Hilbert's school

Formalism: a way of philosophizing about mathematics – older than the *Grundlagenkrise*.

Grundlagenkrise (crisis of foundations, German): a common expression to describe the situation after the paradoxes of set theory.

Formalism, Hilbert's program, Hilbert's school

Formalism: a way of philosophizing about mathematics – older than the *Grundlagenkrise*.

Grundlagenkrise (crisis of foundations, German): a common expression to describe the situation after the paradoxes of set theory.

Formalists say that mathematics is not *about* some ideal objects but is just a game with signs governed by certain rules.

Formalism, Hilbert's program, Hilbert's school

Formalism: a way of philosophizing about mathematics – older than the *Grundlagenkrise*.

Grundlagenkrise (crisis of foundations, German):
a common expression to describe the situation
after the paradoxes of set theory.

Formalists say that mathematics is not *about* some ideal objects but is just a game with signs governed by certain rules.

Hilbert's program: proving the reliability of mathematics in metamathematics.
Metamathematics treats mathematical theories *as* sets of strings structured by derivation rules regardless of of their meaning.

Formalism, Hilbert's program, Hilbert's school

Formalism: a way of philosophizing about mathematics – older than the *Grundlagenkrise*.

Grundlagenkrise (crisis of foundations, German): a common expression to describe the situation after the paradoxes of set theory.

Formalists say that mathematics is not *about* some ideal objects but is just a game with signs governed by certain rules.

Hilbert's program: proving the reliability of mathematics in metamathematics.
Metamathematics treats mathematical theories *as* sets of strings structured by derivation rules regardless of of their meaning.

Hilbert's school: young mathematicians working in the 1920's in Göttingen on this program (Wilhelm Ackermann, Paul Bernays, John von Neumann, Jacques Herbrand).

There will be no class next Friday!
Next class: November 6.