Metatheorems about first-order logic

András Máté

13 Nov. 2026

Last week's key lemma, more exactly

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Every step of the algorithm produces at least one consistent extension of Γ and either

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Every step of the algorithm produces at least one consistent extension of Γ and either

- some Γ^* is a finished analytic sequence for Γ

  or

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Every step of the algorithm produces at least one consistent extension of Γ and either

I some Γ^* is a finished analytic sequence for Γ

or

II every Γ^* is finite and contains a trivial contradiction.

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Every step of the algorithm produces at least one consistent extension of Γ and either

I some Γ^* is a finished analytic sequence for Γ

or

II every Γ^* is finite and contains a trivial contradiction.

In case I, Γ^* has a model (therefore Γ has a model, too) whose domain consists of natural numbers only.

Last week's key lemma, more exactly

There exists an algorithm that produces a finite set of Γ^* sequences of closed sentences from Γ s.t.:

Every step of the algorithm produces at least one consistent extension of Γ and either

I some Γ^* is a finished analytic sequence for Γ
or

II every Γ^* is finite and contains a trivial contradiction.

In case I, Γ^* has a model (therefore Γ has a model, too) whose domain consists of natural numbers only.
In case II, Γ is inconsistent.

Theorems

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

- If every finite subset of Γ is consistent, then Γ is consistent, too.

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

- If every finite subset of Γ is consistent, then Γ is consistent, too.
- If a sentence follows (semantically) from Γ (i.e., $\Gamma \cup \{\neg C\}$ has no model), then it follows from a finite part of Γ.

Theorems

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

- If every finite subset of Γ is consistent, then Γ is consistent, too.
- If a sentence follows (semantically) from Γ (i.e., $\Gamma \cup \{\neg C\}$ has no model), then it follows from a finite part of Γ.

Completeness: Every consistent set of sentences Γ has a model. Alternative formulation:

Theorems

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

- If every finite subset of Γ is consistent, then Γ is consistent, too.
- If a sentence follows (semantically) from Γ (i.e., $\Gamma \cup \{\neg C\}$ has no model), then it follows from a finite part of Γ.

Completeness: Every consistent set of sentences Γ has a model. Alternative formulation:

- If a sentence C is a (semantical) consequence of a set of sentences Γ, then C is derivable from Γ.

Compactness: If Γ is inconsistent, then it has an inconsistent finite subset. In other words:

- If every finite subset of Γ is consistent, then Γ is consistent, too.
- If a sentence follows (semantically) from Γ (i.e., $\Gamma \cup \{\neg C\}$ has no model), then it follows from a finite part of Γ.

Completeness: Every consistent set of sentences Γ has a model. Alternative formulation:

- If a sentence C is a (semantical) consequence of a set of sentences Γ, then C is derivable from Γ.

Löwenheim-Skolem: If a set of sentences has a model, then it has a countable model, too.

Some consequences

Some consequences

1. In first-order logic, there is no sentence that expresses the infinity of the domain.

Some consequences

1. In first-order logic, there is no sentence that expresses the infinity of the domain.

2. Suppose that we have a first-order theory of real numbers which contains the usual operations and relations on real numbers and proves at least some simple propositions about them. Suppose further than we have a model for this theory, consisting of real numbers as we used to thinking about them ('standard model').

Some consequences

1. In first-order logic, there is no sentence that expresses the infinity of the domain.

2. Suppose that we have a first-order theory of real numbers which contains the usual operations and relations on real numbers and proves at least some simple propositions about them. Suppose further than we have a model for this theory, consisting of real numbers as we used to thinking about them ('standard model').

Within the theory we can prove that the domain (the set of real numbers) is not countable (Cantor's theorem).

Some consequences

1. In first-order logic, there is no sentence that expresses the infinity of the domain.

2. Suppose that we have a first-order theory of real numbers which contains the usual operations and relations on real numbers and proves at least some simple propositions about them. Suppose further than we have a model for this theory, consisting of real numbers as we used to thinking about them ('standard model').

Within the theory we can prove that the domain (the set of real numbers) is not countable (Cantor's theorem).

But according to Löwenheim-Skolem, the theory has a model where the domain is countable. (Skolem's paradox.)

Some consequences

1. In first-order logic, there is no sentence that expresses the infinity of the domain.

2. Suppose that we have a first-order theory of real numbers which contains the usual operations and relations on real numbers and proves at least some simple propositions about them. Suppose further than we have a model for this theory, consisting of real numbers as we used to thinking about them ('standard model').

Within the theory we can prove that the domain (the set of real numbers) is not countable (Cantor's theorem).

But according to Löwenheim-Skolem, the theory has a model where the domain is countable. (Skolem's paradox.)

Not a contradiction; but it implies that some important notions (e.g. countability) are incurably relative, model-dependent. (Putnam: 'Models and reality', 1980)

Consequences continued

Consequences continued

3. There are non-standard models of the first-order theory of real numbers, e.g. those whith infinitely small positive numbers.

3. There are non-standard models of the first-order theory of real numbers, e.g. those whith infinitely small positive numbers.

Let us consider the following set of propositions:

$$\{0 < a < 1,\ 0 < a < 1/2,\ \ldots,\ 0 < a < 1/n,\ \ldots\}$$
$$\cup\{\text{Axioms of the theory}\}$$

3. There are non-standard models of the first-order theory of real numbers, e.g. those whith infinitely small positive numbers.

Let us consider the following set of propositions:

$$\{0 < a < 1,\ 0 < a < 1/2,\ \ldots,\ 0 < a < 1/n,\ \ldots\}$$
$$\cup\{\text{Axioms of the theory}\}$$

Every finite subset of this set has a model (namely the standard one extended by an appropriate interpretation of 'a'). Therefore, (due to compactness) the whole set has a model, too, and this is also a model of the axioms.

Consequences finished

Consequences finished

4. Similarly, first-order Peano arithmetics has models with infinitely large numbers.

4. Similarly, first-order Peano arithmetics has models with infinitely large numbers.

BTW., nonstandard models of Peano arithmetics can be characterized by the following 2-order sentence:

$$\exists X(\exists x Xx \wedge \forall x(Xx \to x > 0) \wedge$$
$$\forall y[\forall x(Xx \to x > y) \to \forall x(Xx \to x > y')])$$

Gödel's discovery

Gödel's discovery

Gödel 1931: 'On Formally Undecidable Propositions of Principia Mathematica and Related Systems'

Gödel's discovery

Gödel 1931: 'On Formally Undecidable Propositions of Principia Mathematica and Related Systems'

First Incompleteness Theorem: Peano arithmetics is not negation complete.
There is some sentence G such that neither G itself nor $\neg G$ can be deduced from the axioms
(provided that Peano-arithmetics is ω-consistent).

Gödel's discovery

Gödel 1931: 'On Formally Undecidable Propositions of Principia Mathematica and Related Systems'

First Incompleteness Theorem: Peano arithmetics is not negation complete.
There is some sentence G such that neither G itself nor $\neg G$ can be deduced from the axioms
(provided that Peano-arithmetics is ω-consistent).

The statement of the theorem remains valid if the system is extended with new axioms or axiom schemes.
It also holds for systems in which Peano arithmetics has a model (e.g. set theory).

Gödel's discovery

Gödel 1931: 'On Formally Undecidable Propositions of Principia Mathematica and Related Systems'

First Incompleteness Theorem: Peano arithmetics is not negation complete.
There is some sentence G such that neither G itself nor $\neg G$ can be deduced from the axioms
(provided that Peano-arithmetics is ω-consistent).

The statement of the theorem remains valid if the system is extended with new axioms or axiom schemes.
It also holds for systems in which Peano arithmetics has a model (e.g. set theory).

Rosser 1936: Instead of ω-consistency, consistency is enough.

Gödel's discovery

Gödel 1931: 'On Formally Undecidable Propositions of Principia Mathematica and Related Systems'

First Incompleteness Theorem: Peano arithmetics is not negation complete.
There is some sentence G such that neither G itself nor $\neg G$ can be deduced from the axioms
(provided that Peano-arithmetics is ω-consistent).

The statement of the theorem remains valid if the system is extended with new axioms or axiom schemes.
It also holds for systems in which Peano arithmetics has a model (e.g. set theory).

Rosser 1936: Instead of ω-consistency, consistency is enough.

Second Incompleteness Theorem: The sentence expressing the consistency of Peano arithmetics is neither provable nor refutable (under the same conditions and with the same generalizations).

Kalmár's proof of the first incompleteness theorem

Kalmár's proof of the first incompleteness theorem

Language: first-order logic with 0 as an individual constant and some function symbols for arithmetic operations. Include at least the successor ($'$) and the four basic operations ($+$, $*$, $-$, $\div$).

Kalmár's proof of the first incompleteness theorem

Language: first-order logic with 0 as an individual constant and some function symbols for arithmetic operations. Include at least the successor ($'$) and the four basic operations ($+$, $*$, $-$, $\div$).

Numerals are the individual terms 0, $0'$, $0''$, ...
Numerical terms are the terms containing no variable.

Kalmár's proof of the first incompleteness theorem

Language: first-order logic with 0 as an individual constant and some function symbols for arithmetic operations. Include at least the successor ($'$) and the four basic operations ($+$, $*$, $-$, $\div$).

Numerals are the individual terms $0,\ 0',\ 0'',\ \ldots$.
Numerical terms are the terms containing no variable.
Assume that we can calculate the value of any numerical term.
Calculating a numerical term t means proving some equality $t = n$ (where n is a numeral).

A matrix of inequalities and its diagonal

A matrix of inequalities and its diagonal

Consider the terms of the language that contain (at most) one free variable. These can be enumerated in an (infinite) sequence:

$$k_0(x),\ k_1(x),\ \ldots, k_n(x),\ \ldots$$

The indexes are the Gödel numbers of the terms.

A matrix of inequalities and its diagonal

Consider the terms of the language that contain (at most) one free variable. These can be enumerated in an (infinite) sequence:

$$k_0(x),\ k_1(x),\ \ldots, k_n(x),\ \ldots$$

The indexes are the Gödel numbers of the terms.

Let us arrange the inequalities of the form $k_n(x) \neq m$ in a two-dimensional infinite table in the obvious way:

A matrix of inequalities and its diagonal

Consider the terms of the language that contain (at most) one free variable. These can be enumerated in an (infinite) sequence:

$$k_0(x),\ k_1(x),\ \ldots, k_n(x),\ \ldots$$

The indexes are the Gödel numbers of the terms.

Let us arrange the inequalities of the form $k_n(x) \neq m$ in a two-dimensional infinite table in the obvious way:

$$\begin{array}{llll} k_0(x) \neq 0 & k_0(x) \neq 1 & \ldots & k_0(x) \neq n \quad \ldots \\ k_1(x) \neq 0 & k_1(x) \neq 1 & \ldots & k_1(x) \neq n \quad \ldots \\ \vdots & & & \\ k_n(x) \neq 0 & k_n(x) \neq 1 & \ldots & k_n(x) \neq n \quad \ldots \end{array}$$

Diagonalization

Diagonalization

If we have some effective axiom system and derivation rules (i.e. an effectively axiomatized theory), some of these inequalities become provable, others become refutable. Are there 'neither-nor' cases?

Diagonalization

If we have some effective axiom system and derivation rules (i.e. an effectively axiomatized theory), some of these inequalities become provable, others become refutable. Are there 'neither-nor' cases?

Consider the diagonal of the table, i. e. the sequence of formulas $k_n(x) \neq n$ (call them diagonal formulas). We can enumerate all the proofs in our theory, and therefore we can also enumerate the proofs that prove diagonal formulas:

$$P_0,\ P_1,\ \ldots,\ P_n,\ \ldots$$

Diagonalization

If we have some effective axiom system and derivation rules (i.e. an effectively axiomatized theory), some of these inequalities become provable, others become refutable. Are there 'neither-nor' cases?

Consider the diagonal of the table, i. e. the sequence of formulas $k_n(x) \neq n$ (call them diagonal formulas). We can enumerate all the proofs in our theory, and therefore we can also enumerate the proofs that prove diagonal formulas:

$$P_0,\ P_1,\ \ldots,\ P_n,\ \ldots$$

Define the function f as $f(n) = m$ iff P_n proves the mth diagonal formula.

Diagonalization

If we have some effective axiom system and derivation rules (i.e. an effectively axiomatized theory), some of these inequalities become provable, others become refutable. Are there 'neither-nor' cases?

Consider the diagonal of the table, i. e. the sequence of formulas $k_n(x) \neq n$ (call them diagonal formulas). We can enumerate all the proofs in our theory, and therefore we can also enumerate the proofs that prove diagonal formulas:

$$P_0,\ P_1,\ \ldots,\ P_n,\ \ldots$$

Define the function f as $f(n) = m$ iff P_n proves the mth diagonal formula.

Lemma (not proved): $f(x)$ can be expressed in our language by a term with one variable.

The Gödel sentence

The Gödel sentence

A consequence of the above lemma: there is at least one of the expressions $< k_n(x) >$ which expresses $f(x)$. Let g be the index of the first such expression. I.e., for all x, $f(x) = k_g(x)$

The Gödel sentence

A consequence of the above lemma: there is at least one of the expressions $< k_n(x) >$ which expresses $f(x)$. Let g be the index of the first such expression. I.e., for all x, $f(x) = k_g(x)$

Consider the gth diagonal formula:

$$k_g(x) \neq g \tag{G}$$

The Gödel sentence

A consequence of the above lemma: there is at least one of the expressions $< k_n(x) >$ which expresses $f(x)$. Let g be the index of the first such expression. I.e., for all x, $f(x) = k_g(x)$

Consider the gth diagonal formula:

$$k_g(x) \neq g \tag{G}$$

If (G) is provable, then for some m, the proof P_m proves G, therefore by the definition of f, $f(m) = k_g(m) = g$, and so (G) is false.

The Gödel sentence

A consequence of the above lemma: there is at least one of the expressions $< k_n(x) >$ which expresses $f(x)$. Let g be the index of the first such expression. I.e., for all x, $f(x) = k_g(x)$

Consider the gth diagonal formula:

$$k_g(x) \neq g \tag{G}$$

If (G) is provable, then for some m, the proof P_m proves G, therefore by the definition of f, $f(m) = k_g(m) = g$, and so (G) is false.
If (G) is false, then for some n, $k_g(n) = f(n) = g$, and therefore P_n proves (G).

The Gödel sentence

A consequence of the above lemma: there is at least one of the expressions $< k_n(x) >$ which expresses $f(x)$. Let g be the index of the first such expression. I.e., for all x, $f(x) = k_g(x)$

Consider the gth diagonal formula:

$$k_g(x) \neq g \tag{G}$$

If (G) is provable, then for some m, the proof P_m proves G, therefore by the definition of f, $f(m) = k_g(m) = g$, and so (G) is false.
If (G) is false, then for some n, $k_g(n) = f(n) = g$, and therefore P_n proves (G).
In summary, (G) is provable iff it is false.

The final result and some discussion

The final result and some discussion

If our arithmetic (this could be Peano arithmetic or any effective extension of it) calculates every numerical term and *proves only true equalities with at most one variable*, then the Gödel sentence (G) is true and not provable, and its negation is not provable because it is false. Therefore it is negation incomplete.

The final result and some discussion

If our arithmetic (this could be Peano arithmetic or any effective extension of it) calculates every numerical term and *proves only true equalities with at most one variable*, then the Gödel sentence (G) is true and not provable, and its negation is not provable because it is false. Therefore it is negation incomplete.

A plausible reading of the sentence (G): For every x, the value of the expression $k_g(x)$ (therefore $f(x)$) is different from g. This means that the diagonal formula numbered with g has no proof. But the g-th diagonal formula is (G) itself!! Therefore (G) says: 'I am not provable'.

The final result and some discussion

If our arithmetic (this could be Peano arithmetic or any effective extension of it) calculates every numerical term and *proves only true equalities with at most one variable*, then the Gödel sentence (G) is true and not provable, and its negation is not provable because it is false. Therefore it is negation incomplete.

A plausible reading of the sentence (G): For every x, the value of the expression $k_g(x)$ (therefore $f(x)$) is different from g. This means that the diagonal formula numbered with g has no proof. But the g-th diagonal formula is (G) itself!! Therefore (G) says: 'I am not provable'.

For theories satisfying the italic condition above, we have proved the first incompleteness theorem.

The final result and some discussion

If our arithmetic (this could be Peano arithmetic or any effective extension of it) calculates every numerical term and *proves only true equalities with at most one variable*, then the Gödel sentence (G) is true and not provable, and its negation is not provable because it is false. Therefore it is negation incomplete.

A plausible reading of the sentence (G): For every x, the value of the expression $k_g(x)$ (therefore $f(x)$) is different from g. This means that the diagonal formula numbered with g has no proof. But the g-th diagonal formula is (G) itself!! Therefore (G) says: 'I am not provable'.

For theories satisfying the italic condition above, we have proved the first incompleteness theorem.

Gödel applied a weaker condition than the above: he assumed that that the theory was ω-consistent.

The final result and some discussion

If our arithmetic (this could be Peano arithmetic or any effective extension of it) calculates every numerical term and *proves only true equalities with at most one variable*, then the Gödel sentence (G) is true and not provable, and its negation is not provable because it is false. Therefore it is negation incomplete.

A plausible reading of the sentence (G): For every x, the value of the expression $k_g(x)$ (therefore $f(x)$) is different from g. This means that the diagonal formula numbered with g has no proof. But the g-th diagonal formula is (G) itself!! Therefore (G) says: 'I am not provable'.

For theories satisfying the italic condition above, we have proved the first incompleteness theorem.

Gödel applied a weaker condition than the above: he assumed that that the theory was ω-consistent.

A consistent theory is $\underline{\omega\text{-inconsistent}}$ iff there is some property P s.t. the theory proves $P(0)$, $P(1)$, ... $P(n)$, ... for each numeral n, but it proves $\exists x \neg P(x)$, too.

The (un)provability of consistency

The (un)provability of consistency

The consistency of PA can be expressed within PA by a formula CPA:

The (un)provability of consistency

The consistency of PA can be expressed within PA by a formula CPA:

CPA $\leftrightarrow$ There is no natural number s.t. it is the Gödel number of the proof of $0 = 0'$

The (un)provability of consistency

The consistency of PA can be expressed within PA by a formula CPA:

CPA $\leftrightarrow$ There is no natural number s.t. it is the Gödel number of the proof of $0 = 0'$

CPA is a deductively undecidable sentence in PA. (This is the second incompleteness theorem.) It is true on the standard model but false on some non-standard models.

The (un)provability of consistency

The consistency of PA can be expressed within PA by a formula CPA:

CPA $\leftrightarrow$ There is no natural number s.t. it is the Gödel number of the proof of $0 = 0'$

CPA is a deductively undecidable sentence in PA. (This is the second incompleteness theorem.) It is true on the standard model but false on some non-standard models.

PA + $\neg$ CPA is an example of a consistent, but ω-inconsistent theory (provided that Peano arithmetics is consistent).

Impact of the second incompleteness theorem

- Gödel: 'I wish to note expressly that [this theorem] does not contradict Hilbert's formalistic viewpoint. For this viewpoint presupposes only the existence of a consistency proof in which nothing but finitary means of proof is used and it is conceivable that there exist finitary proofs that *cannot* be expressed in the formalism of [first-order Peano arithmetics].' (Original paper on the incompleteness theorems)

Impact of the second incompleteness theorem

- Gödel: 'I wish to note expressly that [this theorem] does not contradict Hilbert's formalistic viewpoint. For this viewpoint presupposes only the existence of a consistency proof in which nothing but finitary means of proof is used and it is conceivable that there exist finitary proofs that *cannot* be expressed in the formalism of [first-order Peano arithmetics].' (Original paper on the incompleteness theorems)
- von Neumann: 'Thus I am today of the opinion that
  1. Gödel has shown the unrealizability of Hilbert's program.
  2. There is no more reason to reject intuitionism (if one disregards the aesthetic issue, which in practice also for me be the decisive factor).'

  (Letter to Carnap, 1931)